

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: C12N 5/00, 15/63, 15/09, A61K 48/00		A1	(11) International Publication Number: WO 98/01538
			(43) International Publication Date: 15 January 1998 (15.01.98)
(21) International Application Number: PCT/US97/11956		(81) Designated States: AU, CA, IL, JP, KR, MX, NO, NZ, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 9 July 1997 (09.07.97)			
(30) Priority Data: 08/677,762 10 July 1996 (10.07.96) US 08/763,995 12 December 1996 (12.12.96) US		Published With international search report.	
(71) Applicant: IMMUNEX CORPORATION [US/US]; 51 University Street, Seattle, WA 98101 (US).			
(72) Inventors: MARASKOVSKY, Eugene; 4123 Evanston Avenue North, Seattle, WA 98103 (US). MCKENNA, Hilary, R.; 3809 Whitman Avenue North, Seattle, WA 98103 (US).			
(74) Agent: PERKINS, Patricia, Anne; Immunex Corporation, 51 University Street, Seattle, WA 98101 (US).			
(54) Title: METHOD OF ACTIVATING DENDRITIC CELLS			
(57) Abstract <p>Antigen-expressing, activated dendritic cells are disclosed. Such dendritic cells are used to present tumor, viral or bacterial antigens to T cells, and can be useful in vaccination protocols. Other cytokines can be used in separate, sequential or simultaneous combination with the activated, antigen-pulsed dendritic cells. Also disclosed are methods for stimulating an immune response using the antigen-expressing, activated dendritic cells.</p>			

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SE	Sweden
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BD	Bangladesh	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	ME	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BT	Bhutan	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Cote d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroun	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

TITLE
METHOD OF ACTIVATING DENDRITIC CELLS

FIELD OF THE INVENTION

The present invention relates to a dendritic cell activation factor, to methods of enhancing a lymphocyte-mediated immune response in vivo, and to dendritic cell populations useful in the manipulation of cellular and humoral immune responses.

BACKGROUND OF THE INVENTION

Vaccination is an efficient means of preventing death or disability from infectious diseases. The success of this method in the field of infectious disease has also stimulated interest in utilizing vaccination in the treatment or prevention of neoplastic disease. Despite the successes achieved with the use of vaccines, however, there are still many challenges in the field of vaccine development. Parenteral routes of administration, the numbers of different vaccinations required and the need for, and frequency of, booster immunizations all impede efforts to control or eliminate disease.

One such difficulty is lack of immunogenicity in an antigen, i.e., the antigen is unable to promote an effective immune response against the pathogen. In addition, certain antigens may elicit only a certain type of immune response, for example, a cell-mediated or a humoral response. Adjuvants are substances that enhance, augment or potentiate an immune response, and can in some instances be used to promote one type of immune response over another. Although numerous vaccine adjuvants are known, alum is the only adjuvant widely used in humans.

Dendritic cells are a heterogeneous cell population with distinctive morphology and a widespread tissue distribution (Steinman, R.M., *Annu. Rev. Immunol.*, 9:271-296, 1991). Dendritic cells are referred to as "professional" antigen presenting cells, and have a high capacity for sensitizing MHC-restricted T cells. Thus, there is growing interest in using dendritic cells ex vivo as tumor or infectious disease vaccine adjuvants (see, for example, Romani, et al., *J. Exp. Med.*, 180:83, 1994). Therefore, an agent that enhanced the ability of dendritic cells to stimulate an immune response would be of wide importance.

35

SUMMARY OF THE INVENTION

The present invention pertains to a method of activating dendritic cells to enhance antigen presenting capacity. The activated, antigen-presenting dendritic cells of the invention are useful as vaccine adjuvants.

The invention also provides a method of generating large quantities of antigen-presenting dendritic cells ex vivo. Following collection of an individual's CD34⁺ hematopoietic progenitors and stem cells, cytokines such as granulocyte-macrophage colony stimulating factor (GM-CSF) and flt-3 ligand (flt3-L) can be used to expand the cells in vitro and to drive them to differentiate into cells of the dendritic cell lineage. Cytokines can also be used to increase the numbers of CD34⁺ cells in circulation prior to collection. The resulting dendritic cells are exposed to an antigen one wishes to elicit an immune response against, and allowed to process the antigen (this procedure is sometimes referred to in the art as "antigen-pulsing"). The antigen-pulsed (or antigen-expressing) dendritic cells are then activated with a CD40 binding protein, and subsequently administered to the individual.

An alternate method for preparing dendritic cells that present antigen is to transfect the dendritic cells with a gene encoding an antigen or a specific polypeptide derived therefrom. Once the dendritic cells express the antigen in the context of MHC, the dendritic cells are activated with a CD40 binding protein, and subsequently administered to the individual to provide a stronger and improved immune response to the antigen.

The activated antigen-presenting dendritic cells can also be used as a vaccine adjuvant and can be administered prior to, concurrently with or subsequent to antigen administration. Moreover, the dendritic cells can be administered to the individual prior to, concurrently with or subsequent to administration of cytokines that modulate an immune response, for example a CD40 binding protein (i.e., soluble CD40L), or a soluble CD83 molecule. Additional useful cytokines include, but are not limited to, Interleukins (IL) 1, 2, 4, 5, 6, 7, 10, 12 and 15, colony stimulating factors (CSF) such as GM-CSF, granulocyte colony stimulating factor (G-CSF), or GM-CSF/IL-3 fusion proteins, or other cytokines such as TNF- α or c-kit ligand. Moreover, biologically active derivatives of these cytokines; and combinations thereof will also be useful.

The invention also provides for the ex vivo preparation of antigen-specific T cells. Following the procedures described above for preparing large numbers of antigen-presenting dendritic cells ex vivo, the collected antigen-presenting dendritic cells are used to generate antigen-specific T cells from naive T cells that have been collected from the individual. After the antigen has been adequately presented to the T cells generated, the antigen-specific T cells can be administered to the individual.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 presents the results of an allo-antigen T cell proliferation assay, demonstrating that incubation of dendritic cells with CD40L prior to their use in an MLR

(mixed lymphocyte reaction) increases the ability of the dendritic cells to stimulate the proliferation of T cells by about threefold, as described in Example 3.

Figure 2 illustrates that dendritic cells that are cultured with CD40L are less effective at presenting antigen to antigen-specific T cells than dendritic cells that were not exposed to CD40L, as described in Example 4.

Figure 3 demonstrates that dendritic cells that are first pulsed with antigen, then cultured with CD40L are more effective at presenting antigen to antigen-specific T cells than dendritic cells that were pulsed with antigen but not exposed to CD40L, as described in Example 5.

DETAILED DESCRIPTION OF THE INVENTION

The invention is directed to the use of CD40L to activate antigen-pulsed dendritic cells. Activation enhances the ability of the dendritic cells to present antigen to lymphoid cells, and thus augments the immune response against the antigen. Another embodiment of the invention is the isolation and use of activated, antigen-pulsed dendritic cells as vaccine adjuvants. The activated, antigen-pulsed dendritic cells may also be used *ex vivo* to generate antigen-specific T cells.

Dendritic cells

Dendritic cells comprise a heterogeneous cell population with distinctive morphology and a widespread tissue distribution. The dendritic cell system and its role in immunity is reviewed by Steinman, R.M., *Annu. Rev. Immunol.*, 9:271-296 (1991), incorporated herein by reference. The cell surface of dendritic cells is unusual, with characteristic veil-like projections, and is characterized by having the cell surface markers CD1a⁺, CD4⁺, CD86⁺, or HLA-DR⁺. Dendritic cells have a high capacity for sensitizing MHC-restricted T cells and are very effective at presenting antigens to T cells *in situ*, both self-antigens during T cell development and tolerance and foreign antigens during immunity.

Because of their effectiveness at antigen presentation, there is growing interest in using dendritic cells *ex vivo* as tumor or infectious disease vaccine adjuvants (see, for example, Romani, et al., *J. Exp. Med.*, 180:83 (1994)). The use of dendritic cells as immunostimulatory agents has been limited due to the low frequency of dendritic cells in peripheral blood, the limited accessibility of lymphoid organs and the dendritic cells' terminal state of differentiation. Dendritic cells originate from CD34⁺ bone marrow or peripheral blood progenitors and peripheral blood mononuclear cells, and the proliferation and maturation of dendritic cells can be enhanced by the cytokines GM-CSF (sargramostim, Leukine®, Immunex Corporation, Seattle, Washington), TNF- α , c-kit

ligand (also known as stem cell factor (SCF), steel factor (SF), or mast cell growth factor (MGF)) and interleukin-4. Recently, flt3-L has been found to stimulate the generation of large numbers of functionally mature dendritic cells, both in vivo and in vitro (USSN 08/539,142, filed October 4, 1995).

5

Ex vivo culture of dendritic cells

A procedure for ex vivo expansion of hematopoietic stem and progenitor cells is described in U.S. Patent No. 5,199,942, incorporated herein by reference. Other suitable methods are known in the art. Briefly, ex vivo culture and expansion comprises: (1) collecting CD34⁺ hematopoietic stem and progenitor cells from a patient from peripheral blood harvest or bone marrow explants; and (2) expanding such cells *ex vivo*. In addition to the cellular growth factors described in Patent 5,199,942, other factors such as flt3-L, IL-1, IL-3 and c-kit ligand, can be used.

Stem or progenitor cells having the CD34 marker constitute only about 1% to 3% of the mononuclear cells in the bone marrow. The amount of CD34⁺ stem or progenitor cells in the peripheral blood is approximately 10- to 100-fold less than in bone marrow. Cytokines such as flt3-L may be used to increase or mobilize the numbers of dendritic cells in vivo. Increasing the quantity of an individual's dendritic cells may facilitate antigen presentation to T cells for antigen(s) that already exists within the patient, such as a tumor antigen, or a bacterial or viral antigen. Alternatively, cytokines may be administered prior to, concurrently with or subsequent to administration of an antigen to an individual for immunization purposes.

Peripheral blood cells are collected using apheresis procedures known in the art. See, for example, Bishop et al., *Blood*, vol. 83, No. 2, pp. 610-616 (1994). Briefly, peripheral blood progenitor cells (PBPC) and peripheral blood stem cells (PBSC) are collected using conventional devices, for example, a Haemonetics Model V50 apheresis device (Haemonetics, Braintree, MA). Four-hour collections are performed typically no more than five times weekly until approximately 6.5×10^8 mononuclear cells (MNC)/kg are collected. The cells are suspended in standard media and then centrifuged to remove red blood cells and neutrophils. Cells located at the interface between the two phases (the buffy coat) are withdrawn and resuspended in HBSS. The suspended cells are predominantly mononuclear and a substantial portion of the cell mixture are early stem cells.

A variety of cell selection techniques are known for identifying and separating CD34⁺ hematopoietic stem or progenitor cells from a population of cells. For example, monoclonal antibodies (or other specific cell binding proteins) can be used to bind to a marker protein or surface antigen protein found on stem or progenitor cells. Several such markers or cell surface antigens for hematopoietic stem cells (i.e., flt-3, CD34, My-10, and

Thy-1) are known in the art, as are specific binding proteins therefore (see for example, U.S.S.N. 08/539, 142, filed October 4, 1995).

In one method, antibodies or binding proteins are fixed to a surface, for example, glass beads or flask, magnetic beads, or a suitable chromatography resin, and contacted with the population of cells. The stem cells are then bound to the bead matrix. Alternatively, the binding proteins can be incubated with the cell mixture and the resulting combination contacted with a surface having an affinity for the antibody-cell complex. Undesired cells and cell matter are removed providing a relatively pure population of stem cells. The specific cell binding proteins can also be labeled with a fluorescent label, e.g., chromophore or fluorophore, and the labeled cells separated by sorting. Preferably, isolation is accomplished by an immunoaffinity column.

Immunoaffinity columns can take any form, but usually comprise a packed bed reactor. The packed bed in these bioreactors is preferably made of a porous material having a substantially uniform coating of a substrate. The porous material, which provides a high surface area-to-volume ratio, allows for the cell mixture to flow over a large contact area while not impeding the flow of cells out of the bed. The substrate should, either by its own properties, or by the addition of a chemical moiety, display high-affinity for a moiety found on the cell-binding protein. Typical substrates include avidin and streptavidin, while other conventional substrates can be used.

In one useful method, monoclonal antibodies that recognize a cell surface antigen on the cells to be separated are typically further modified to present a biotin moiety. The affinity of biotin for avidin thereby removably secures the monoclonal antibody to the surface of a packed bed (see Berenson, et al., *J. Immunol. Meth.*, 91:11, 1986). The packed bed is washed to remove unbound material, and target cells are released using conventional methods. Immunoaffinity columns of the type described above that utilize biotinylated anti-CD34 monoclonal antibodies secured to an avidin-coated packed bed are described for example, in WO 93/08268.

An alternative means of selecting the quiescent stem cells is to induce cell death in the dividing, more lineage-committed, cell types using an antimetabolite such as 5-fluorouracil (5-FU) or an alkylating agent such as 4-hydroxycyclophosphamide (4-HC). The non-quiescent cells are stimulated to proliferate and differentiate by the addition of growth factors that have little or no effect on the stem cells, causing the non-stem cells to proliferate and differentiate and making them more vulnerable to the cytotoxic effects of 5-FU or 4-HC. See Berardi et al., *Science*, 267:104 (1995), which is incorporated herein by reference.

Isolated stem cells can be frozen in a controlled rate freezer (e.g., Cryo-Med, Mt. Clemens, MI), then stored in the vapor phase of liquid nitrogen using dimethylsulfoxide as

a cryoprotectant. A variety of growth and culture media can be used for the growth and culture of dendritic cells (fresh or frozen), including serum-depleted or serum-based media. Useful growth media include RPMI, TC 199, Iscove's modified Dulbecco's medium (Iscove, et al., *F.J. Exp. Med.*, 147:923 (1978)), DMEM, Fischer's, alpha medium, NCTC, F-10, Leibovitz's L-15, MEM and McCoy's.

Particular nutrients present in the media include serum albumin, transferrin, lipids, cholesterol, a reducing agent such as 2-mercaptoethanol or monothioglycerol, pyruvate, butyrate, and a glucocorticoid such as hydrocortisone 2-hemisuccinate. More particularly, the standard media includes an energy source, vitamins or other cell-supporting organic compounds, a buffer such as HEPES, or Tris, that acts to stabilize the pH of the media, and various inorganic salts. A variety of serum-free cellular growth media is described in WO 95/00632, which is incorporated herein by reference.

The collected CD34⁺ cells are cultured with suitable cytokines, for example, as described herein, and in USSN 08/539,142. CD34⁺ cells then are allowed to differentiate and commit to cells of the dendritic lineage. These cells are then further purified by flow cytometry or similar means, using markers characteristic of dendritic cells, such as CD1a, HLA DR, CD80 and/or CD86. The cultured dendritic cells are exposed to an antigen, for example, a tumor antigen or an antigen derived from a pathogenic or opportunistic organism, allowed to process the antigen, and then cultured with an amount of a CD40 binding protein to activate the dendritic cell. Alternatively, the dendritic cells are transfected with a gene encoding an antigen, and then cultured with an amount of a CD40 binding protein to activate the antigen-presenting dendritic cells.

The activated, antigen-carrying dendritic cells are then administered to an individual in order to stimulate an antigen-specific immune response. The dendritic cells can be administered prior to, concurrently with, or subsequent to, antigen administration. Alternatively, T cells may be collected from the individual and exposed to the activated, antigen-carrying dendritic cells in vitro to stimulate antigen-specific T cells, which are administered to the individual.

30 Useful cytokines

Various cytokines will be useful in the ex vivo culture of dendritic cells. Flt3-L refers to a genus of polypeptides that are described in EP 0627487 A2 and in WO 94/28391, both incorporated herein by reference. A human flt3-L cDNA was deposited with the American Type Culture Collection, Rockville, Maryland, USA (ATCC) on August 6, 1993 and assigned accession number ATCC 69382. IL-3 refers to a genus of interleukin-3 polypeptides as described in U.S. Patent No. 5,108,910, incorporated herein by reference. A DNA sequence encoding human IL-3 protein suitable for use in the

invention is publicly available from the American Type Culture Collection (ATCC) under accession number ATCC 67747. c-kit ligand is also referred to as Mast Cell Growth Factor (MGF), Steel Factor or Stem Cell Factor (SCF), and is described in EP 423,980, which is incorporated herein by reference.

- 5 Other useful cytokines include Interleukin-4 (IL-4; Mosley et al., *Cell* 59:335 (1989), Idzerda et al., *J. Exp. Med.* 171:861 (1990) and Galizzi et al., *Int. Immunol.* 2:669 (1990), each of which is incorporated herein by reference) and granulocyte-macrophage colony stimulating factor (GM-CSF; described in U.S. Patent Nos. 5,108,910, and 5,229,496 each of which is incorporated herein by reference).
10 Commercially available GM-CSF (sargramostim, Leukine®) is obtainable from Immunex Corp., Seattle, WA). Moreover, GM-CSF/IL-3 fusion proteins (i.e., a C-terminal to N-terminal fusion of GM-CSF and IL-3) will also be useful in ex vivo culture of dendritic cells. Such fusion proteins are known and are described in U.S. Patent Nos. 5,199,942, 5,108,910 and 5,073,627, each of which is incorporated herein by reference. A preferred
15 fusion protein is PIXY321 as described in US Patent No. 5,199,942.

- In addition to their use in ex vivo culture of dendritic cells, cytokines will also be useful in the present invention by separate, sequential or simultaneous administration of a cytokine or cytokines with activated, antigen-pulsed dendritic cells. Preferred cytokines are those that modulate an immune response, particularly cytokines selected from the group
20 consisting of Interleukins 1, 2, 3, 4, 5, 6, 7, 10, 12 and 15; granulocyte-macrophage colony stimulating factor, granulocyte colony stimulating factor, a fusion protein comprising Interleukin-3 and granulocyte-macrophage colony stimulating factor, Interferon- γ , TNF, TGF- β , flt-3 ligand; soluble CD40 ligand; biologically active derivatives of these cytokines; and combinations thereof. Soluble CD83, described in USSN 08/601,954, filed
25 February 15, 1996), and soluble CD40L (described in USSN 08/477,733 and USSN 08/484,624, both filed June 7, 1995) are particularly preferred cytokines.

- Useful cytokines act by binding a receptor present on the surface of a dendritic cell and transducing a signal. Moreover, additional binding proteins can be prepared as described herein for CD40 binding proteins, that bind appropriate cytokine receptors and
30 transduce a signal to a dendritic cell. For example, WO 95/27062 describes agonistic antibodies to Flt-3, the receptor for Flt-3L, from which various Flt-3 binding proteins can be prepared. Additional useful cytokines include biologically active analogs of cytokines that are useful for culturing dendritic cells. Useful cytokine analogs have an amino acid sequence that is substantially similar to the native cytokine, and are biologically active
35 capable of binding to their specific receptor and transducing a biological signal. Such analogs can be prepared and tested by methods that are known in the art and as described herein.

CD40/CD40L

CD40 is a member of the tumor necrosis factor (TNF)/nerve growth factor (NGF) receptor family, which is defined by the presence of cysteine-rich motifs in the extracellular region (Smith et al., *Science* 248:1019, 1990; Mallett and Barclay, *Immunology Today* 12:220, 1991). This family includes the lymphocyte antigen CD27, CD30 (an antigen found on Hodgkin's lymphoma and Reed-Sternberg cells), two receptors for TNF, a murine protein referred to as 4-1BB, rat OX40 antigen, NGF receptor, and Fas antigen. Human CD40 antigen (CD40) is a peptide of 277 amino acids having a molecular weight of 30,600 (Stamenkovic et al., *EMBO J.* 8:1403, 1989).

Activated CD4+ T cells express high levels of a ligand for CD40 (CD40L). Human CD40L was cloned from peripheral blood T-cells as described in Spriggs et al., *J. Exp. Med.* 176:1543 (1992). The cloning of murine CD40L is described in Armitage et al., *Nature* 357:80 (1992). CD40L is a type II membrane polypeptide having an extracellular region at its C-terminus, a transmembrane region and an intracellular region at its N-terminus. CD40L biological activity is mediated by binding of the extracellular region of CD40L with CD40, and includes B cell proliferation and induction of antibody secretion (including IgE secretion).

CD40L is believed to be important in feedback regulation of an immune response. For example, a CD40+ antigen presenting cell will present antigen to a T cell, which will then become activated and express CD40L. The CD40L will, in turn, further activate the antigen presenting cell, increasing its efficiency at antigen presentation, and upregulating expression of Class I and Class II MHC, CD80 and CD86 costimulatory molecules, as well as various cytokines (Caux et al., *J. Exp. Med.* 180:1263, 1994).

Useful forms of CD40L for the inventive methods as disclosed in USSN 08/477,733 and USSN 08/484,624, both filed June 7, 1995, and both of which are incorporated by reference herein. Such useful forms include soluble, oligomeric CD40 ligand comprising a CD40-binding peptide and an oligomer-forming peptide. The CD40-binding peptide is selected from the group consisting of:

- (a) a peptide comprising amino acids 1 through 261, 35 through 261, 34 through 225, 113 through 261, 113 through 225, 120 through 261, or 120 through 225 of SEQ ID NO:2;
- (b) fragments of a peptide according to (a) that bind CD40; and
- (c) peptides encoded by DNA which hybridizes to a DNA that encodes a peptide of (a) or (b), under stringent conditions (hybridization in 6 X SSC at 63°C overnight; washing in 3 X SSC at 55°C), and which bind to CD40.

Useful oligomer-forming peptides are also disclosed in USSN 08/477,733 and USSN 08/484,624, and exemplified in SEQ ID NOs: 3 and 4 herein.

- A corresponding family of ligands exists for molecules in the TNFR family, and several of these are also expressed on activated T cells or other cells of the immune system.
- 5 This family includes tumor necrosis factor and lymphotoxin (TNF and LT, respectively; reviewed in Ware et al., *Curr. Top. Microbiol. Immunol.* 198:175, 1995), as well as CD27L (USSN 08/106,507, filed August 13, 1993), CD30L (US Patent 5,480,981, issued January 2, 1996), 4-1BBL (USSN 08/236,918, filed May 6, 1994), OX40L (US patent 5,457,035, issued October 10, 1995) and Fas L (USSN 08/571,579, filed
- 10 December 13, 1995). These ligands are also known to be involved in modulation of an immune response, and are likely to be useful to activate antigen-pulsed dendritic cells or other antigen presenting cells that bear the corresponding receptor.

CD40 Monoclonal Antibodies and Additional CD40 Binding Proteins

- 15 Useful CD40 binding proteins are those that are capable of binding CD40 and inhibiting binding of CD40 to CD40L, as determined by observing at least about 90% inhibition of the binding of soluble CD40 to CD40L, and include monoclonal antibodies, CD40 ligand, and molecules derived therefrom. Monoclonal antibodies directed against the CD40 surface antigen (CD40 mAb) have been shown to mediate various biological
- 20 activities on human B cells (see for example, LEUKOCYTE TYPING IV; A.J. McMichael ed. Oxford University Press, Oxford, p. 426). U.S.S.N. 08/130, 541, filed October 1, 1993, the relevant disclosure of which is incorporated by reference, discloses two monoclonal antibodies that specifically bind CD40, referred to as hCD40m2 and hCD40m3. Unlike other CD40 mAb, hCD40m2 (ATCC HB11459; deposited under terms of the Budapest
- 25 Treaty with the American Type Culture Collection in Rockville, MD, USA, on October 6, 1993) and hCD40m3 bind CD40 and inhibit binding of CD40 to cells that constitutively express CD40L, indicating that hCD40m2 and hCD40m3 bind CD40 in or near the ligand binding domain.

- Additional CD40 monoclonal antibodies may be generated using conventional
- 30 techniques (see U.S. Patent Nos. RE 32,011, 4,902,614, 4,543,439, and 4,411,993 which are incorporated herein by reference; see also *Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses*, Plenum Press, Kennett, McKearn, and Bechtol (eds.), 1980, and *Antibodies: A Laboratory Manual*, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, 1988, which are also incorporated herein by reference).
- 35 Monoclonal antibodies that bind CD40 in or near the ligand binding domain will also be useful in the present invention.

Additional CD40 binding proteins may also be constructed utilizing recombinant DNA techniques. For example, the variable regions of a gene which encodes an antibody to CD40 that binds in or near the ligand binding domain can be incorporated into a useful CD40 binding protein (see Larrick et al., *Biotechnology* 7:934, 1989; Reichmann et al., *Nature* 332:323, 1988; Roberts et al., *Nature* 328:731, 1987; Verhoeyen et al., *Science* 239:1534, 1988; Chaudhary et al., *Nature* 339:394, 1989).

Briefly, DNA encoding the antigen-binding site (or CD40 binding domain; variable region) of a CD40 mAb is isolated, amplified, and linked to DNA encoding another protein, for example a human IgG (see Verhoeyen et al., *supra*; see also Reichmann et al., *supra*). Alternatively, the antigen-binding site (variable region) may be either linked to, or inserted into, another completely different protein (see Chaudhary et al., *supra*), resulting in a new protein with antigen-binding sites of the antibody as well as the functional activity of the completely different protein.

Similarly, the CD40 binding region (extracellular domain) of a CD40 ligand may be used to prepare other CD40 binding proteins. Useful forms of CD40 ligand are disclosed in U.S.S.N. 08/477,733 and U.S.S.N. 08/484,624, both of which were filed on June 7, 1995. Additional forms of CD40 ligand can be prepared by methods known in the art. As for other useful CD40 binding proteins, CD40 ligand will bind CD40 in or near the ligand binding domain, and will be capable of transducing a signal to a cell expressing CD40 (i.e., biologically active).

DNA sequences that encode proteins or peptides that form oligomers will be particularly useful in preparation of CD40 binding proteins comprising an antigen binding domain of CD40 antibody, or an extracellular domain of a CD40 ligand. Certain of such oligomer-forming proteins are disclosed in U.S.S.N. 08/477,733 and U.S.S.N. 08/484,624, both of which were filed on June 7, 1995; additional, useful oligomer-forming proteins are also disclosed in U.S.S.N. 08/446,922, filed May 18, 1995. Fc fusion proteins (including those that are formed with Fc muclins have decreased affinity for Fc receptors) can also be prepared.

Mutant forms of CD40 binding proteins that are substantially similar (i.e., those having an amino acid sequence at least 80% identical to a native amino acid sequence, most preferably at least 90% identical) to the previously described CD40 binding proteins will also be useful in the present invention. The percent identity may be determined, for example, by comparing sequence information using the GAP computer program, version 6.0 described by Devereux et al. (*Nucl. Acids Res.* 12:387, 1984) and available from the University of Wisconsin Genetics Computer Group (UWGCG). The GAP program utilizes the alignment method of Needleman and Wunsch (*J. Mol. Biol.* 48:443, 1970), as revised by Smith and Waterman (*Adv. Appl. Math* 2:482, 1981). The preferred default

parameters for the GAP program include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) for nucleotides, and the weighted comparison matrix of Gribskov and Burgess, *Nucl. Acids Res.* 14:6745, 1986, as described by Schwartz and Dayhoff, eds., *Atlas of Protein Sequence and Structure*,
5 National Biomedical Research Foundation, pp. 353-358, 1979; (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps.

Generally, substitutions of different amino acids from those in the native form of a useful CD40 binding protein should be made conservatively; i.e., the most preferred
10 substitute amino acids are those which do not affect the ability of the inventive proteins to bind CD40 in a manner substantially equivalent to that of native CD40 ligand. Examples of conservative substitutions include substitution of amino acids outside of the binding domain(s), and substitution of amino acids that do not alter the secondary and/or tertiary structure of CD40 binding proteins. Additional examples include substituting one aliphatic
15 residue for another, such as Ile, Val, Leu, or Ala for one another, or substitutions of one polar residue for another, such as between Lys and Arg; Glu and Asp; or Gln and Asn. Other such conservative substitutions, for example, substitutions of entire regions having similar hydrophobicity characteristics, are well known.

Similarly, when a deletion or insertion strategy is adopted, the potential effect of the
20 deletion or insertion on biological activity should be considered. Subunits of CD40 binding proteins may be constructed by deleting terminal or internal residues or sequences. Additional guidance as to the types of mutations that can be made is provided by a comparison of the sequence of CD40 binding proteins to proteins that have similar structures

25 Mutations must, of course, preserve the reading frame phase of the coding sequences and preferably will not create complementary regions that could hybridize to produce secondary mRNA structures such as loops or hairpins which would adversely affect translation of the CD40 binding protein mRNA. Although a mutation site may be predetermined, it is not necessary that the nature of the mutation *per se* be predetermined.
30 For example, in order to select for optimum characteristics of mutants at a given site, random mutagenesis may be conducted at the target codon and the expressed mutated proteins screened for the desired activity.

Mutations can be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of
35 the native sequence. Following ligation, the resulting reconstructed sequence encodes an analog having the desired amino acid insertion, substitution, or deletion. Alternatively, oligonucleotide-directed site-specific mutagenesis procedures can be employed to provide

an altered gene having particular codons altered according to the substitution, deletion, or insertion required. Exemplary methods of making the alterations set forth above are disclosed by Walder et al. (*Gene* 42:133, 1986); Bauer et al. (*Gene* 37:73, 1985); Craik (*BioTechniques*, January 1985, 12-19); Smith et al. (*Genetic Engineering: Principles and Methods*, Plenum Press, 1981); and U.S. Patent Nos. 4,518,584 and 4,737,462 disclose suitable techniques, and are incorporated by reference herein.

As is well-known in the art, not all mutations will cause a change in amino acid sequence. Mutations that confer advantageous properties in the production of recombinant proteins will also be useful for preparing useful CD40 binding proteins. Naturally occurring variants are also encompassed by the invention. Examples of such variants are proteins that result from alternate mRNA splicing events or from proteolytic cleavage of the protein, wherein the native biological property is retained.

Once suitable antibodies or binding proteins have been obtained, they may be isolated or purified by many techniques well known to those of ordinary skill in the art (see *Antibodies: A Laboratory Manual*, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, 1988). Suitable techniques include peptide or protein affinity columns, HPLC or RP-HPLC, purification on protein A or protein G columns, or any combination of these techniques. Recombinant CD40 binding proteins can be prepared according to standard methods, and tested for binding specificity to CD40 utilizing assays known in the art, including for example ELISA, ABC, or dot blot assays, as well by bioactivity assays. The latter will also be useful in evaluating the biological activity of CD40 binding proteins.

Preparation of antigens

Immunization is a centuries old, and highly effective, means of inducing a protective immune response against pathogens in order to prevent or ameliorate disease. The vaccines that have been used for such induction are generally live, attenuated microorganisms, or preparations of killed organisms or fractions thereof. Live, attenuated vaccines are generally thought to more closely mimic the immune response that occurs with a natural infection than do those prepared from killed microbes or non-infective preparations derived from pathogens (i.e., toxoids, recombinant protein vaccines). However, attenuated vaccines also present a risk of reversion to pathogenicity, and can cause illness, especially in immunocompromised individuals.

Along with improved sanitation, immunization has been the most efficient means of preventing death or disability from numerous infectious diseases in humans and in other animals. Vaccination of susceptible populations has been responsible for eliminating small pox world wide, and for drastic decreases in the occurrence of such diseases as diphtheria, pertussis, and paralytic polio in the developed nations. Numerous vaccines are licensed for

administration to humans, including live virus vaccines for certain adenoviruses, measles, mumps and rubella viruses, and poliovirus, diphtheria and tetanus toxoid vaccines, and *Haemophilus b* and meningococcal polysaccharide vaccines (Hinman et al., in Principles and Practice of Infectious Diseases, 3rd edition; G.L. Mandell, R.G. Douglas and J.E. Bennett, eds., Churchill Livingstone Inc., NY, NY; 2320-2333; Table 2).

5 In addition to use in the area of infectious disease, vaccination is also considered a promising therapy for cancer. For such uses, tumor-associated antigens can be prepared from tumor cells, either by preparing crude lysates of tumor cells, for example as described in Cohen et al., *Cancer Res.* 54:1055 (1994) and Cohen et al., *Eur. J. Immunol.* 24:315
10 (1994), or by partially purifying the antigens (for example, as described by Itoh et al., *J. Immunol.* 153:1202; 1994). Moreover, useful tumor antigens may be purified further, or even expressed recombinantly, to provide suitable antigen preparations. Any other methods of identifying and isolating antigens against which an immune response would be beneficial in cancer will also find utility in the inventive methods.

15 Purified dendritic cells are then pulsed with (exposed to) antigen, to allow them to take up the antigen in a manner suitable for presentation to other cells of the immune systems. Antigens are classically processed and presented through two pathways. Peptides derived from proteins in the cytosolic compartment are presented in the context of Class I MHC molecules, whereas peptides derived from proteins that are found in the
20 endocytic pathway are presented in the context of Class II MHC. However, those of skill in the art recognize that there are exceptions; for example, the response of CD8⁺ tumor specific T cells, which recognize exogenous tumor antigens expressed on MHC Class I. A review of MHC-dependent antigen processing and peptide presentation is found in Germain, R.N., *Cell* 76:287 (1994).

25 Numerous methods of pulsing dendritic cells with antigen are known; those of skill in the art regard development of suitable methods for a selected antigen as routine experimentation. In general, the antigen is added to cultured dendritic cells under conditions promoting viability of the cells, and the cells are then allowed sufficient time to take up and process the antigen, and express antigen peptides on the cell surface in
30 association with either Class I or Class II MHC, a period of about 24 hours (from about 18 to about 30 hours, preferably 24 hours). Dendritic cells may also be exposed to antigen by transfecting them with DNA encoding the antigen. The DNA is expressed, and the antigen is presumably processed via the cytosolic/Class I pathway.

35 Administration of activated, antigen-pulsed dendritic cells

The present invention provides methods of using therapeutic compositions comprising activated, antigen-pulsed dendritic cells. The use of such cells in conjunction

with soluble cytokine receptors or cytokines, or other immunoregulatory molecules is also contemplated. The inventive compositions are administered to stimulate an immune response, and can be given by bolus injection, continuous infusion, sustained release from implants, or other suitable technique. Typically, the cells on the inventive methods will be administered in the form of a composition comprising the antigen-pulsed, activated dendritic cells in conjunction with physiologically acceptable carriers, excipients or diluents. Such carriers will be nontoxic to recipients at the dosages and concentrations employed. Neutral buffered saline or saline mixed with conspecific serum albumin are exemplary appropriate diluents.

For use in stimulating a certain type of immune response, administration of other cytokines along with activated, antigen-pulsed dendritic cells is also contemplated. Several useful cytokines (or peptide regulatory factors) are discussed in Schrader, J.W. (*Mol Immunol* 28:295; 1991). Such factors include (alone or in combination) Interleukins 1, 2, 4, 5, 6, 7, 10, 12 and 15; granulocyte-macrophage colony stimulating factor, granulocyte colony stimulating factor; a fusion protein comprising Interleukin-3 and granulocyte-macrophage colony stimulating factor; Interferon- γ , TNF, TGF- β , flt-3 ligand and biologically active derivatives thereof. A particularly preferred cytokine is CD40 ligand (CD40L). A soluble form of CD40L is described in USSN 08/484,624, filed June 7, 1995. Other cytokines will also be useful, as described herein. DNA encoding such cytokines will also be useful in the inventive methods, for example, by transfecting the dendritic cells to express the cytokines. Administration of these immunomodulatory molecules includes simultaneous, separate or sequential administration with the cells of the present invention.

The relevant disclosures of all publications cited herein are specifically incorporated by reference. The following examples are provided to illustrate particular embodiments and not to limit the scope of the invention.

EXAMPLE 1

This Example describes a method for generating purified dendritic cells *ex vivo*. Human bone marrow is obtained, and cells having a CD34⁺ phenotype are isolated using a CD34 antibody column (CellPro, Bothell, WA). The CD34⁺ cells are cultured in a suitable medium, for example, McCoy's enhanced media, that contains cytokines that promote the growth of dendritic cells (i.e., 20 ng/ml each of GM-CSF, IL-4, TNF- α , or 100 ng/ml flt3-L or c-kit ligand, or combinations thereof). The culture is continued for approximately two weeks at 37 °C in 10% CO₂ in humid air. Cells then are sorted by flow cytometry using antibodies for CD1a⁺, HLA-DR⁺ and CD86⁺. A combination of GM-CSF, IL-4 and

TNF- α can yield a six to seven-fold increase in the number of cells obtained after two weeks of culture, of which 50-80% of cells are CD1a⁺ HLA-DR⁺ CD86⁺. The addition of flt3-L and/or c-kit ligand further enhances the expansion of total cells, and therefore of the dendritic cells. Phenotypic analysis of cells isolated and cultured under these conditions indicates that between 60-70% of the cells are HLA-DR⁺, CD86⁺, with 40-50% of the cells expressing CD1a in all factor combinations examined.

EXAMPLE 2

This Example describes a method for collecting and expanding dendritic cells. Prior to cell collection, flt3-L or sargramostim (Leukine[®], Immunex Corporation, Seattle, WA) may be administered to an individual to mobilize or increase the numbers of circulating PBPC and PBSC. Other growth factors such as CSF-1, GM-CSF, c-kit ligand, G-CSF, EPO, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, GM-CSF/IL-3 fusion proteins, LIF, FGF and combinations thereof, can be likewise administered in sequence, or in concurrent combination with flt3-L.

Mobilized or non-mobilized PBPC and PBSC are collected using apheresis procedures known in the art. See, for example, Bishop et al., *Blood*, vol. 83, No. 2, pp. 610-616 (1994). Briefly, PBPC and PBSC are collected using conventional devices, for example, a Haemonetics Model V50 apheresis device (Haemonetics, Braintree, MA). Four-hour collections are performed typically no more than five times weekly until approximately 6.5×10^8 mononuclear cells (MNC)/kg individual are collected.

Aliquots of collected PBPC and PBSC are assayed for granulocyte-macrophage colony-forming unit (CFU-GM) content. Briefly, MNC (approximately 300,000) are isolated, cultured at 37 °C in 5% CO₂ in fully humidified air for about two weeks in modified McCoy's 5A medium, 0.3% agar, 200 U/ml recombinant human GM-CSF, 200 u/ml recombinant human IL-3, and 200 u/ml recombinant human G-CSF. Other cytokines, including flt3-L or GM-CSF/IL-3 fusion molecules (PIXY 321), may be added to the cultures. These cultures are stained with Wright's stain, and CFU-GM colonies are scored using a dissecting microscope (Ward et al., *Exp. Hematol.*, 16:358 (1988)). Alternatively, CFU-GM colonies can be assayed using the CD34/CD33 flow cytometry method of Siena et al., *Blood*, Vol. 77, No. 2, pp 400-409 (1991), or any other method known in the art.

CFU-GM containing cultures are frozen in a controlled rate freezer (e.g., Cryo-Med, Mt. Clemens, MI), then stored in the vapor phase of liquid nitrogen. Ten percent dimethylsulfoxide can be used as a cryoprotectant. After all collections from the individual have been made, CFU-GM containing cultures are thawed and pooled, then contacted with flt3-L either alone, sequentially or in concurrent combination with other cytokines listed

above to drive the CFU-GM to dendritic cell lineage. The dendritic cells are cultured and analyzed for percentage of cells displaying selected markers as described above.

EXAMPLE 3

5 This example illustrates the ability of CD40L-stimulated dendritic cells to present allo-antigen and therefore cause proliferation of T cells. CD34+ cells were obtained from the bone marrow of a human donor, cultured for two weeks in the presence of selected cytokines, and isolated by flow cytometry substantially as described in Example 1. Prior to their use in a mixed lymphocyte reaction (MLR), the dendritic cells were cultured for an additional 24 hours in the presence or absence of a soluble trimeric form of CD40L (1 µg/ml) in McCoy's enhanced media containing cytokines that support the growth of dendritic cells.

10 T cells were purified from the blood of a non-HLA matched donor by rosetting with 2-aminoethylisothiuronium bromide hydrobromide-treated sheep red blood cells. CD4+ and CD8+ populations were further purified using immunomagnetic selection using a MACS (Milenyi Biotec, Sunnyvale, CA) according to the manufacturer's protocol. Cell proliferation assays were conducted with the purified T cells in RPMI (10% heat-inactivated fetal bovine serum (FBS)), in the presence of titrated numbers of the dendritic cells, at 37°C in a 10% CO₂ atmosphere. Approximately 1×10^5 T cells per well were cultured in triplicate in round-bottomed 96-well microtiter plates (Corning) for seven days, in the presence of varying numbers of the unmatched dendritic cells. The cells were pulsed with 1 µCi/well of tritiated thymidine (25 Ci/nmole, Amersham, Arlington Heights, IL) for the final eight hours of culture.

20 Cells were harvested onto glass fiber discs with an automated cell harvester and incorporated cpm were measured by liquid scintillation spectrometry. The results, which are shown in Figure 1, demonstrated that three-fold fewer CD40L-activated dendritic cells were required to stimulate the equivalent proliferation of T cells compared to dendritic cells that had not been exposed to CD40L prior to their use in an MLR. This increase was most likely due to increased expression of cell surface molecules that stimulate allo-reactive T cells.

EXAMPLE 4

30 This example illustrates the ability of dendritic cells to stimulate antigen-specific proliferation of T cells. CD34+ cells were obtained from the bone marrow of a human donor believed to be reactive against tetanus toxoid, cultured for two weeks in the presence of selected cytokines, and isolated by flow cytometry substantially as described in Example 1. Prior to their use in a tetanus toxoid (TTX) antigen presentation assay, the dendritic cells were cultured for an additional 24 hours in the presence or absence of a soluble trimeric

form of CD40L (1 µg/ml) in McCoy's enhanced media containing cytokines that support the growth of dendritic cells, then pulsed with purified TTX (Connaught Laboratory Inc., Swiftwater, PA), at 37°C in a 10% CO₂ atmosphere for 24 hrs.

Autologous tetanus toxoid-reactive T cells were derived by culturing the CD34⁺ cells that were eluted from the CD34 antibody column in the presence of purified TTX and low concentrations of IL-2 and IL-7 (2 ng/ml and 5 ng/ml, respectively) for two weeks. The CD34⁺ population contains a percentage of T cells (about 5%), a proportion of which are reactive against tetanus toxoid, as well as other cell types that act as antigen presenting cells. By week 2, analysis of these cells indicated that they were about 90% T cells, the majority of which were tetanus toxoid-specific, with low levels of the T cell activation markers.

Antigen specific T cell proliferation assays were conducted with TTX-specific T cells from CD34⁺ bone marrow cells as above, in RPMI with added 10% heat-inactivated fetal bovine serum (FBS), in the presence of the tetanus toxoid-pulsed dendritic cells, at 37°C in a 10% CO₂ atmosphere. Approximately 1×10^5 T cells per well were cultured in triplicate in round-bottomed 96-well microtiter plates (Corning) for five days, in the presence of a titrated number of dendritic cells. The cells were pulsed with 1 µCi/well of tritiated thymidine (25 Ci/nmole, Amersham, Arlington Heights, IL) for the final four to eight hours of culture. Cells were harvested onto glass fiber discs with an automated cell harvester and incorporated cpm were measured by liquid scintillation spectrometry. The results, which are shown in Figure 2, indicated that dendritic cells that are cultured with CD40L are about ten-fold less efficient at presenting antigen to TTX-specific T cells than dendritic cells that were not exposed to CD40L.

EXAMPLE 5

This example illustrates the ability of CD40L to activate antigen-pulsed dendritic cells for stimulation of antigen-specific T cells. CD34⁺ cells were obtained and treated as described in Example 4, except that the cells were pulsed with tetanus toxoid for 24 hours prior to culture with CD40L. Autologous tetanus toxoid-reactive T cells were derived, and antigen specific T cell proliferation assays conducted, as described in Example 4. The results, which are shown in Figure 3, indicated that three-fold fewer dendritic cells that are first pulsed with antigen, then cultured with CD40L, were required to stimulate the equivalent level of proliferation when presenting TTX to TTX-specific T cells than dendritic cells that were pulsed with antigen but not exposed to CD40L.

SEQUENCE LISTING

- 5 (1) GENERAL INFORMATION:
- (i) APPLICANT: Immunex Corporation
- (ii) TITLE OF INVENTION: METHOD OF ACTIVATING DENDRITIC CELLS
- 10 (iii) NUMBER OF SEQUENCES: 4
- (iv) CORRESPONDENCE ADDRESS:
- 15 (A) ADDRESSEE: IMMUNEX CORPORATION
(B) STREET: 51 UNIVERSITY STREET
(C) CITY: SEATTLE
(D) STATE: WASHINGTON
(E) COUNTRY: USA
(F) ZIP: 98101
- 20 (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: Power Macintosh 7200/90
(C) OPERATING SYSTEM: Apple Operating System 7.6
25 (D) SOFTWARE: Microsoft Word for Macintosh, Version #6.0.1
- (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER: --to be assigned--
(B) FILING DATE: 09-JUL-1997
30 (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: USSN 08/677,762
(B) FILING DATE: 10-JUL-1996
35 (C) CLASSIFICATION:
- (viii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: USSN 08/763,995
(B) FILING DATE: 12-DEC-1996
40 (C) CLASSIFICATION:
- (ix) ATTORNEY/AGENT INFORMATION:
- (A) NAME: Perkins, Patricia A.
(B) REGISTRATION NUMBER: 34,693
45 (C) REFERENCE/DOCKET NUMBER: 2845-WO
- (ix) TELECOMMUNICATION INFORMATION:
- (A) TELEPHONE: (206)587-0430
(B) TELEFAX: (206)233-0644
- 50 (2) INFORMATION FOR SEQ ID NO:1:
- (i) SEQUENCE CHARACTERISTICS:
- 55 (A) LENGTH: 840 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
- 60 (ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

5 (vi) ORIGINAL SOURCE:
(A) ORGANISM: *Homo sapiens*

(vii) IMMEDIATE SOURCE:
(B) CLONE: CD40-L

10 (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 46..831

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

	TGCCACCTTC TCTGCCAGAA GATACCATTT CAACTTTAAC ACAGC ATG ATC GAA	54
		Met Ile Glu
		1
20	ACA TAC AAC CAA ACT TCT CCC CGA TCT GCG GCC ACT GGA CTG CCC ATC	102
	Thr Tyr Asn Gln Thr Ser Pro Arg Ser Ala Ala Thr Gly Leu Pro Ile	
	5 10 15	
25	AGC ATG AAA ATT TTT ATG TAT TTA CTT ACT GTT TTT CTT ATC ACC CAG	150
	Ser Met Lys Ile Phe Met Tyr Leu Leu Thr Val Phe Leu Ile Thr Gln	
	20 25 30 35	
30	ATG ATT GGG TCA GCA CTT TTT GCT GTG TAT CTT CAT AGA AGG TTG GAC	198
	Met Ile Gly Ser Ala Leu Phe Ala Val Tyr Leu His Arg Arg Leu Asp	
	40 45 50	
35	AAG ATA GAA GAT GAA AGG AAT CTT CAT GAA GAT TTT GTA TTC ATG AAA	246
	Lys Ile Glu Asp Glu Arg Asn Leu His Glu Asp Phe Val Phe Met Lys	
	55 60 65	
40	ACG ATA CAG AGA TGC AAC ACA GGA GAA AGA TCC TTA TCC TTA CTG AAC	294
	Thr Ile Gln Arg Cys Asn Thr Gly Glu Arg Ser Leu Ser Leu Leu Asn	
	70 75 80	
45	TGT GAG GAG ATT AAA AGC CAG TTT GAA GGC TTT GTG AAG GAT ATA ATG	342
	Cys Glu Glu Ile Lys Ser Gln Phe Glu Gly Phe Val Lys Asp Ile Met	
	85 90 95	
50	TTA AAC AAA GAG GAG ACG AAG AAA GAA AAC AGC TTT GAA ATG CAA AAA	390
	Leu Asn Lys Glu Glu Thr Lys Lys Glu Asn Ser Phe Glu Met Gln Lys	
	100 105 110 115	
55	GGT GAT CAG AAT CCT CAA ATT GCG GCA CAT GTC ATA AGT GAG GCC AGC	438
	Gly Asp Gln Asn Pro Gln Ile Ala Ala His Val Ile Ser Glu Ala Ser	
	120 125 130	
60	AGT AAA ACA ACA TCT GTG TTA CAG TGG GCT GAA AAA GGA TAC TAC ACC	486
	Ser Lys Thr Thr Ser Val Leu Gln Trp Ala Glu Lys Gly Tyr Tyr Thr	
	135 140 145	
65	ATG AGC AAC AAC TTG GTA ACC CTG GAA AAT GGG AAA CAG CTG ACC GTT	534
	Met Ser Asn Asn Leu Val Thr Leu Glu Asn Gly Lys Gln Leu Thr Val	
	150 155 160	

	AAA AGA CAA GGA CTC TAT TAT ATC TAT GCC CAA GTC ACC TTC TGT TCC	582
	Lys Arg Gln Gly Leu Tyr Tyr Ile Tyr Ala Gln Val Thr Phe Cys Ser	
	165 170 175	
5	AAT CGG GAA GCT TCG AGT CAA GCT CCA TTT ATA GCC AGC CTC TGC CTA	630
	Asn Arg Glu Ala Ser Ser Gln Ala Pro Phe Ile Ala Ser Leu Cys Leu	
	180 185 190 195	
10	AAG TCC CCC GGT AGA TTC GAG AGA ATC TTA CTC AGA GCT GCA AAT ACC	678
	Lys Ser Pro Gly Arg Phe Glu Arg Ile Leu Leu Arg Ala Ala Asn Thr	
	200 205 210	
15	CAC AGT TCC GCC AAA CCT TGC GGG CAA CAA TCC ATT CAC TTG GGA GGA	726
	His Ser Ser Ala Lys Pro Cys Gly Gln Gln Ser Ile His Leu Gly Gly	
	215 220 225	
20	GTA TTT GAA TTG CAA CCA GGT GCT TCG GTG TTT GTC AAT GTG ACT GAT	774
	Val Phe Glu Leu Gln Pro Gly Ala Ser Val Phe Val Asn Val Thr Asp	
	230 235 240	
	CCA AGC CAA GTG AGC CAT GGC ACT GGC TTC ACG TCC TTT GGC TTA CTC	822
	Pro Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe Gly Leu Leu	
	245 250 255	
25	AAA CTC TGAACAGTGT CA	840
	Lys Leu	
	260	
30	(2) INFORMATION FOR SEQ ID NO:2:	
	(i) SEQUENCE CHARACTERISTICS:	
	(A) LENGTH: 261 amino acids	
	(B) TYPE: amino acid	
	(D) TOPOLOGY: linear	
35	(ii) MOLECULE TYPE: protein	
	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:	
40	Met Ile Glu Thr Tyr Asn Gln Thr Ser Pro Arg Ser Ala Ala Thr Gly	
	1 5 10 15	
45	Leu Pro Ile Ser Met Lys Ile Phe Met Tyr Leu Leu Thr Val Phe Leu	
	20 25 30	
	Ile Thr Gln Met Ile Gly Ser Ala Leu Phe Ala Val Tyr Leu His Arg	
	35 40 45	
50	Arg Leu Asp Lys Ile Glu Asp Glu Arg Asn Leu His Glu Asp Phe Val	
	50 55 60	
55	Phe Met Lys Thr Ile Gln Arg Cys Asn Thr Gly Glu Arg Ser Leu Ser	
	65 70 75 80	
	Leu Leu Asn Cys Glu Glu Ile Lys Ser Gln Phe Glu Gly Phe Val Lys	
	85 90 95	
60	Asp Ile Met Leu Asn Lys Glu Glu Thr Lys Lys Glu Asn Ser Phe Glu	
	100 105 110	

Met Gln Lys Gly Asp Gln Asn Pro Gln Ile Ala Ala His Val Ile Ser
 115 120 125

5 Glu Ala Ser Ser Lys Thr Thr Ser Val Leu Gln Trp Ala Glu Lys Gly
 130 135 140

Tyr Tyr Thr Met Ser Asn Asn Leu Val Thr Leu Glu Asn Gly Lys Gln
 145 150 155 160

10 Leu Thr Val Lys Arg Gln Gly Leu Tyr Tyr Ile Tyr Ala Gln Val Thr
 165 170 175

Phe Cys Ser Asn Arg Glu Ala Ser Ser Gln Ala Pro Phe Ile Ala Ser
 180 185 190

15 Leu Cys Leu Lys Ser Pro Gly Arg Phe Glu Arg Ile Leu Leu Arg Ala
 195 200 205

20 Ala Asn Thr His Ser Ser Ala Lys Pro Cys Gly Gln Gln Ser Ile His
 210 215 220

Leu Gly Gly Val Phe Glu Leu Gln Pro Gly Ala Ser Val Phe Val Asn
 225 230 235 240

25 Val Thr Asp Pro Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe
 245 250 255

30 Gly Leu Leu Lys Leu
 260

(2) INFORMATION FOR SEQ ID NO:3:

35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 740 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

45 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: HUMAN

50 (vii) IMMEDIATE SOURCE:
 (B) CLONE: IgG1 Fc

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

55 CGGTACCGCT AGCGTCGACA GGCCTAGGAT ATCGATACGT ACAGCCCAGA TCTTCTGACA 60
 AAACCTCACAC ATGCCACCG TCCCCAGCAC CTGAACCTCT GGGGGGACCG TCAGTCTTCC 120
 60 TCTTCCCCC AAAACCCAAG GACACCTCA TGATCTCCCG GACCCCTGAG GTCACATGCG 180

TGGTGGTGGG CGTGAGCCAC GAAGACCCCTG AGGTCAAGTT CAACTGGTAC GTGGACGGCG 240
 TGGAGGTGCA TAATGCCAAG ACAAGCCCGC GGGAGGAGCA GTACAACAGC ACGTACCCGG 300
 5 TGGTCAGCGT CCTCACCGTC CTGCACCAGG ACTGGCTGAA TGGCAAGGAC TACAAGTGCA 360
 AGGTCTCCAA CAAAGCCCTC CCAGCCCCCA TGCAGAAAAC CATCTCCAAA GCCAAAGGGC 420
 10 AGCCCCGAGA ACCACAGGTG TACACCCTGC CCCCATCCCG GGATGAGCTG ACCAAGAACC 480
 AGGTACAGCT GACCTGCCCTG GTCAAAGGCT TCTATCCAG GCACATCGCC GTGGAGTGGG 540
 AGAGCAATGG GCAGCCGGAG AACAACTACA AGACCACGCC TCCCGTGCTG GACTCCGACG 600
 15 GCTCTTCTT CCTCTACAGC AAGCTCACC TGGACAAGAG CAGGTGGCAG CAGGGGAACG 660
 TCTTCTCATG CTCGGTGATG CATGAGGCTC TGCACAACCA CTACACGCAG AAGAGCCTCT 720
 20 CCCTGTCTCC GGGTAAATGA 740

(2) INFORMATION FOR SEQ ID NO:4:

25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 30 (ii) MOLECULE TYPE: peptide
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:
 35 Arg Met Lys Gln Ile Glu Asp Lys Ile Glu Glu Ile Leu Ser Lys Ile
 1 5 10 15
 Tyr His Ile Glu Asn Glu Ile Ala Arg Ile Lys Lys Leu Ile Gly Glu
 20 25 30
 40 Arg

CLAIMS

What is claimed is:

1. A population of activated dendritic cells that express an antigen, produced by the process of
 - (a) obtaining dendritic cells;
 - (b) causing the dendritic cells to express the antigen by either (i) exposing the dendritic cells to the antigen in culture under conditions promoting uptake and processing of the antigen, or (ii) transfecting the dendritic cells with a gene encoding the antigen; and
 - (c) activating the antigen-expressing dendritic cells by exposing them to a CD40 binding protein capable of binding CD40 and inhibiting binding of CD40 to CD40L, as determined by observing at least about 90% inhibition of the binding of soluble CD40 to CD40L.
2. The population according to claim 1 wherein dendritic cells are obtained by contacting hematopoietic stem or progenitor cells with a molecule selected from the group consisting of GM-CSF, flt-3L, IL-4, TNF- α , IL-3, c-kit ligand, fusions of GM-CSF and IL-3, and combinations thereof.
3. The population according to claim 1, wherein the CD40 binding protein is a soluble, oligomeric CD40 ligand selected from the group consisting of:
 - (a) a peptide comprising amino acids 1 through 261, 35 through 261, 34 through 225, 113 through 261, 113 through 225, 120 through 261, or 120 through 225 of SEQ ID NO:2;
 - (b) fragments of a peptide according to (a) that bind CD40; and
 - (c) peptides encoded by DNA which hybridizes to a DNA that encodes a peptide of (a) or (b), under stringent conditions (hybridization in 6 X SSC at 63°C overnight; washing in 3 X SSC at 55°C), and which bind to CD40, and an oligomer-forming peptide.
4. The population according to claim 3, wherein the soluble, oligomeric CD40 ligand is selected from the group consisting of:
 - (a) a polypeptide having an amino acid sequence as set forth in SEQ ID NO:2 wherein a cysteine at amino acid 194 is replaced with another amino acid; and
 - (b) a polypeptide that is a fragment of the mutein (a) that binds CD40; wherein the amino acid that is substituted for the cysteine at amino acid 194 is selected from the group consisting of tryptophan, serine, aspartic acid, and lysine.

5. A method of stimulating an immune response specific for an antigen in an individual, comprising the steps of:
- (a) obtaining dendritic cells from the individual;
 - (b) causing the dendritic cells to express the antigen by either (i) exposing the dendritic cells to the antigen in culture under conditions promoting uptake and processing of the antigen, or (ii) transfecting the dendritic cells with a gene encoding the antigen;
 - (c) activating the antigen-expressing dendritic cells by exposing them to a CD40 binding protein capable of binding CD40 and inhibiting binding of CD40 to CD40L, as determined by observing at least about 90% inhibition of the binding of soluble CD40 to CD40L; and
 - (d) administering the activated, antigen-expressing dendritic cells to the individual.
6. The method according to claim 5, wherein the dendritic cells are obtained by obtaining hematopoietic stem or progenitor cells from the individual, and contacting the hematopoietic stem or progenitor cells with a molecule selected from the group consisting of flt-3 ligand, GM-CSF, IL-4, TNF- α , IL-3, c-kit ligand, fusions of GM-CSF and IL-3, and combinations thereof.
7. The method according to claim 6, wherein the CD40 binding protein is a soluble, oligomeric CD40 ligand selected from the group consisting of:
- (a) a peptide comprising amino acids 1 through 261, 35 through 261, 34 through 225, 113 through 261, 113 through 225, 120 through 261, or 120 through 225 of SEQ ID NO:2;
 - (b) fragments of a peptide according to (a) that bind CD40; and
 - (c) peptides encoded by DNA which hybridizes to a DNA that encodes a peptide of (a) or (b), under stringent conditions (hybridization in 6 X SSC at 63°C overnight; washing in 3 X SSC at 55°C), and which bind to CD40, and an oligomer-forming peptide.
8. The population according to claim 7, wherein the soluble, oligomeric CD40 ligand is selected from the group consisting of:
- (a) a polypeptide having an amino acid sequence as set forth in SEQ ID NO:2 wherein a cysteine at amino acid 194 is replaced with another amino acid; and
 - (b) a polypeptide that is a fragment of the polypeptide (a) that binds CD40; wherein the amino acid that is substituted for the cysteine at amino acid 194 is selected from the group consisting of tryptophan, serine, aspartic acid, and lysine.

9. The method according to claim 5, wherein flt-3 ligand is administered to the individual prior to obtaining the dendritic cells, to expand the number of progenitor cells in the circulation of the individual.

10. The method according to claim 9, wherein the dendritic cells are obtained by obtaining hematopoietic stem or progenitor cells from the individual, and contacting the hematopoietic stem or progenitor cells with a molecule selected from the group consisting of flt-3 ligand, GM-CSF, IL-4, TNF- α , IL-3, c-kit ligand, fusions of GM-CSF and IL-3, and combinations thereof.

11. The method according to claim 5, wherein the antigen-expressing, activated dendritic cells are administered simultaneously, sequentially or separately with a molecule selected from the group consisting of Interleukins 1, 2, 3, 4, 5, 6, 7, 10, 12 and 15; granulocyte-macrophage colony stimulating factor, granulocyte colony stimulating factor; a fusion protein comprising Interleukin-3 and granulocyte-macrophage colony stimulating factor; Interferon- γ , TNF; TGF- β ; flt-3 ligand; soluble CD40 ligand; soluble CD83; biologically active derivatives of these cytokines; and combinations thereof.

12. The method according to claim 9, wherein the CD40 binding protein is a soluble, oligomeric CD40 ligand selected from the group consisting of:

(a) a peptide comprising amino acids 1 through 261, 35 through 261, 34 through 225, 113 through 261, 113 through 225, 120 through 261, or 120 through 225 of SEQ ID NO:2;

(b) fragments of a peptide according to (a) that bind CD40;

(c) peptides encoded by DNA which hybridizes to a DNA that encodes a peptide of (a) or (b), under stringent conditions (hybridization in 6 X SSC at 63°C overnight; washing in 3 X SSC at 55°C), and which bind to CD40;

(d) a polypeptide according to (a) wherein acysteine at amino acid 194 is replaced with another amino acid selected from the group consisting of tryptophan, serine, aspartic acid, and lysine; and

(e) a fragment of the polypeptide of (d) which binds CD40; and an oligomer-forming peptide.

13. The method according to claim 10, wherein the CD40 binding protein is a soluble, oligomeric CD40 ligand selected from the group consisting of:

(a) a peptide comprising amino acids 1 through 261, 35 through 261, 34 through 225, 113 through 261, 113 through 225, 120 through 261, or 120 through 225 of SEQ ID NO:2;

(b) fragments of a peptide according to (a) that bind CD40;

(c) peptides encoded by DNA which hybridizes to a DNA that encodes a peptide of (a) or (b), under stringent conditions (hybridization in 6 X SSC at 63°C overnight; washing in 3 X SSC at 55°C), and which bind to CD40;

(d) a polypeptide according to (a) wherein acysteine at amino acid 194 is replaced with another amino acid selected from the group consisting of tryptophan, serine, aspartic acid, and lysine; and

(e) a fragment of the polypeptide of (d) which binds CD40;
and an oligomer-forming peptide.

14. The method according to claim 11, wherein the CD40 binding protein is a soluble, oligomeric CD40 ligand selected from the group consisting of:

(a) a peptide comprising amino acids 1 through 261, 35 through 261, 34 through 225, 113 through 261, 113 through 225, 120 through 261, or 120 through 225 of SEQ ID NO:2;

(b) fragments of a peptide according to (a) that bind CD40;

(c) peptides encoded by DNA which hybridizes to a DNA that encodes a peptide of (a) or (b), under stringent conditions (hybridization in 6 X SSC at 63°C overnight; washing in 3 X SSC at 55°C), and which bind to CD40;

(d) a polypeptide according to (a) wherein acysteine at amino acid 194 is replaced with another amino acid selected from the group consisting of tryptophan, serine, aspartic acid, and lysine; and

(e) a fragment of the polypeptide of (d) which binds CD40;
and an oligomer-forming peptide.

15. A method of preparing antigen-specific T cells from an individual comprising the steps of:

(a) obtaining dendritic cells from the individual;

(b) causing the dendritic cells to express the antigen by either (i) exposing the dendritic cells to the antigen in culture under conditions promoting uptake and processing of the antigen, or (ii) transfecting the dendritic cells with a gene encoding the antigen;

(c) activating the antigen-expressing dendritic cells by exposing them to a CD40 binding protein capable of binding CD40 and inhibiting binding of CD40 to CD40L, as

determined by observing at least about 90% inhibition of the binding of soluble CD40 to CD40L; and.

(d) allowing the dendritic cells to present the antigen to T cells.

16. The method according to claim 15, wherein the antigen-specific T cells are obtained from the individual, exposed to the antigen-presenting dendritic cells *ex vivo*, and re-administered to the individual.

17. The method according to claim 15, wherein the CD40 binding protein is a soluble, oligomeric CD40 ligand selected from the group consisting of:

(a) a peptide comprising amino acids 1 through 261, 35 through 261, 34 through 225, 113 through 261, 113 through 225, 120 through 261, or 120 through 225 of SEQ ID NO:2;

(b) fragments of a peptide according to (a) that bind CD40;

(c) peptides encoded by DNA which hybridizes to a DNA that encodes a peptide of (a) or (b), under stringent conditions (hybridization in 6 X SSC at 63°C overnight; washing in 3 X SSC at 55°C), and which bind to CD40;

(d) a polypeptide according to (a) wherein acysteine at amino acid 194 is replaced with another amino acid selected from the group consisting of tryptophan, serine, aspartic acid, and lysine; and

(e) a fragment of the polypeptide of (d) which binds CD40;

and an oligomer-forming peptide.

18. The method according to claim 16, wherein the CD40 binding protein is a soluble, oligomeric CD40 ligand selected from the group consisting of:

(a) a peptide comprising amino acids 1 through 261, 35 through 261, 34 through 225, 113 through 261, 113 through 225, 120 through 261, or 120 through 225 of SEQ ID NO:2;

(b) fragments of a peptide according to (a) that bind CD40;


(c) peptides encoded by DNA which hybridizes to a DNA that encodes a peptide of (a) or (b), under stringent conditions (hybridization in 6 X SSC at 63°C overnight; washing in 3 X SSC at 55°C), and which bind to CD40;


(d) a polypeptide according to (a) wherein acysteine at amino acid 194 is replaced with another amino acid selected from the group consisting of tryptophan, serine, aspartic acid, and lysine; and

(e) a fragment of the polypeptide of (d) which binds CD40;


and an oligomer-forming peptide.

19. A population of antigen-specific T cells produced by the process of
- (a) obtaining dendritic cells from an individual;
 - (b) causing the dendritic cells to express the antigen by either (i) exposing the dendritic cells to the antigen in culture under conditions promoting uptake and processing of the antigen, or (ii) transfecting the dendritic cells with a gene encoding the antigen;
 - (c) activating the antigen-expressing dendritic cells by exposing them to a CD40 binding protein capable of binding CD40 and inhibiting binding of CD40 to CD40L, as determined by observing at least about 90% inhibition of the binding of soluble CD40 to CD40L; and
 - (d) allowing the dendritic cells to present the antigen to T cells.
20. The population according to claim 19, wherein the CD40 binding protein is a soluble, oligomeric CD40 ligand selected from the group consisting of:
- (a) a peptide comprising amino acids 1 through 261, 35 through 261, 34 through 225, 113 through 261, 113 through 225, 120 through 261, or 120 through 225 of SEQ ID NO:2;
 - (b) fragments of a peptide according to (a) that bind CD40;
 - (c) peptides encoded by DNA which hybridizes to a DNA that encodes a peptide of (a) or (b), under stringent conditions (hybridization in 6 X SSC at 63°C overnight; washing in 3 X SSC at 55°C), and which bind to CD40;
 - (d) a polypeptide according to (a) wherein a cysteine at amino acid 194 is replaced with another amino acid selected from the group consisting of tryptophan, serine, aspartic acid, and lysine; and
 - (e) a fragment of the polypeptide of (d) which binds CD40;
- and an oligomer-forming peptide.

Allo-antigen T cell proliferation**FIGURE 1**

TTX-specific T cell proliferation**FIGURE 2**

SUBSTITUTE SHEET (RULE 26)

**FIGURE 3**

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/11936

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) : C12N 5/00, 15/63, 15/09; A61K 48/00;

US CL : A35/375, 172.3, 69.1, 6; 424/93.21

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/375, 172.3, 69.1, 6; 424/93.21

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN, MEDLINE, BIOSIS

search terms: dendritic, cd40, transfection, apc

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	PEGUET-NAVARRO ET AL. FUNCTIONAL EXPRESSION OF CD40 ANTIGEN ON HUMAN EPIDERMAL LANGERHANS CELLS. J. OF IMMUNOLOGY. 01 NOVEMBER 1995, VOL. 155, No. 9, PAGES 4241-4247.	1-20
Y	VAN DER BERG ET AL. PROPERTIES OF MOUSE CD40: DIFFERENTIAL EXPRESSION OF CD40 EPITOPES ON DENDRITIC CELLS AND EPITHELIAL CELLS. IMMUNOLOGY. 06 JUNE 1996, VOL. 88, No. 2, PAGES 294-300.	1-20
Y	GROUARD ET AL. HUMAN FOLLICULAR DENDRITIC CELLS ENHANCE CYTOKINE-DEPENDENT GROWTH AND DIFFERENTIATION OF CD40-ACTIVATED B CELLS. J. OF IMMUNOLOGY. 01 OCTOBER 1995, VOL. 155, No. 7, PAGES 3345-3352.	1-20

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	* ¹ later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* ^A document defining the general state of the art which is not considered to be of particular relevance	* ² document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* ^B earlier document published on or after the international filing date	* ³ documents of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* ^C document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* ⁴ document member of the same patent family
* ^D document referring to an oral disclosure, use, exhibition or other means	
* ^E document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

09 SEPTEMBER 1997

Date of mailing of the international search report

15 OCT 1997

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231
Facsimile No. (703) 305-3230

Authorized officer

KAREN M. HALL

Telephone No.

(703) 308-0196

Form PCT/ISA/210 (second sheet) (July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/11956

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y, P	KOCH et al. High Level IL-12 Production by Murine Dendritic Cells: Upregulation via MHC Class II and CD40 Molecules and Downregulation by IL-4 and IL-10. J. of Experimental Medicine. 01 August 1996, Vol. 184, No. 2, pages 741-746.	1-20
Y	JAISWAL et al. Regulation of CD40 ligand expression on naive CD4 T cells: a role for TCR but not co-stimulatory signals. International Immunology. February 1996, Vol. 8, No. 2, pages 275-285.	1-20
Y	GIROLOMONI et al. Immortal mouse dendritic cell lines: New models to investigate the immunobiology of dendritic cells. European J. of Dermatology. January 1995, Vol. 5, No. 1, pages 12-15.	1-20
Y	LEYVA-COBIAN et al. Participation of intracellular oxidative pathways in antigen processing by dendritic cells, B cells and macrophages. Immunology Letters. 1994, Vol. 43, pages 29-37.	1-20
Y	PAGLIA et al. Immortalized Dendritic Cell Line Fully Competent in Antigen Presentation Initiates Primary T Cell Responses In Vivo. J. Experimental Medicine. December 1993, Vol. 178, pages 1893-1901.	1-20

This Page Blank (uspto)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

This Page Blank (uspto)